

Product Name: EuroChem FEATHERTOP 520 Herbicide
APVMA Approval No: 86193/114954



Label Name:	EuroChem FEATHERTOP 520 Herbicide
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	ACTIVE CONSTITUENT: : 520 g/L HALOXYFOP present as the haloxyfop-P methyl ester
Mode of Action:	GROUP A HERBICIDE
Statement of Claims:	For the post-emergent control of a wide range of annual and perennial grass weeds in grain legume and oilseed crops, lucerne, medic and clover pasture and seed crops, forestry, bananas, citrus, grapes, pineapples, pome and stone fruit, pyrethrum, tropical fruit and nut crops as specified in the Directions For Use.
Net Contents:	Contents: 250 mL, 1L, 5L, 10L, 20L
Restraints:	DO NOT apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (waterlogged or drought affected), poor nutrition or previous herbicide treatment as reduced levels of control may result. DO NOT spray if rain is likely to occur within one hour.
Directions for Use:	
Other Limitations:	PROTECTION OF LIVESTOCK DO NOT graze or cut treated crops for stock food except as specified under withholding periods.

	PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT EuroChem FEATHERTOP 520 Herbicide is toxic to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used container.
Withholding Periods:	HARVESTING WITHHOLDING PERIODS Canola, chickpeas, cotton, cowpea, faba beans, field peas, lentils, linola, linseed, lupins, mung beans, navy beans, orchard crops, peanuts, plantation crops, soybeans, sunflowers, vetch or vine crops. NOT REQUIRED WHEN USED AS DIRECTED Medic and Clover Seed Crops: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION STOCK FOOD WITHHOLDING PERIODS: Medic and Clover Pasture: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION Lucerne: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION Canola, Chickpeas, Cotton, Cowpea, Faba Beans, Field Peas, Lentils, Linola, Linseed, Lupins, Mung Beans, Navy Beans, Peanuts, Soybeans, Sunflowers, Vetch: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION COTTON GIN TRASH MUST NOT BE FED TO ANIMALS.
Trade Advice:	
General Instructions:	
Resistance Warning:	GROUP A HERBICIDE EuroChem FEATHERTOP 520 Herbicide is a member of the aryloxyphenoxy propionate group of herbicides. The product has the acetyl CoA carboxylase inhibitor mode of action. For weed resistance management EuroChem FEATHERTOP 520 Herbicide is a Group A herbicide. Some naturally occurring weed biotypes resistant to the product and other inhibitors of acetyl CoA carboxylase herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by the product or other inhibitors of acetyl CoA carboxylase. . Since the occurrence of resistant weeds is difficult to detect prior to use, EuroChem Pty Ltd accepts no liability for any losses that may result from the failure of the product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local EuroChem Pty Ltd representative.
Precautions:	
Protections:	PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS - EuroChem FEATHERTOP 520 Herbicide damages cereals and grasses. - DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. - Cereal crops or grasses planted within twelve weeks of application may be damaged

	by the residual effects of EuroChem FEATHERTOP 520 Herbicide, particularly on light and red soils. PROTECTION OF LIVESTOCK DO NOT graze or cut treated crops for stock food except as specified under withholding periods. PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT EuroChem FEATHERTOP 520 Herbicide is toxic to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used container.
--	--

Storage and Disposal:	Store in the closed, original container, in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT store near feedstuffs, fertilisers or seeds. Triple or preferably pressure rinse containers before disposal. Add rinsate to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product. SMALL SPILL MANAGEMENT Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, cat litter or clay granules to the spill. When absorption is complete, sweep up material and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary wash the spill area with an alkali detergent and water and absorb this wash liquid for disposal as described above.
-----------------------	---

Safety Directions:	Harmful if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing spray and using the prepared spray, wear cotton overalls over normal clothing buttoned to the neck and wrist, a washable hat, elbow- length chemical resistant gloves and disposable face mask covering mouth and nose. For aerial spraying equipment wear cotton overalls buttoned to the neck and wrist and elbow- length chemical resistant gloves. Wash hands after each day's use, wash gloves and contaminated clothing.
--------------------	---

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 1126. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor
-------------------------	--

First Aid Warnings:	
---------------------	--

DIRECTIONS FOR USE**Table 1a. Winter crops - Canola, Chickpeas, Faba beans, Field peas, Lentils, Linola, Linseed, Lupins, Lucerne, Vetch, Medic and Clover pastures or seed crops.**

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE (mL/ha)		CRITICAL COMMENTS
		With Uptake Spraying Oil	With a Non-ionic Wetter	
Annual ryegrass	2 to 4 leaf	75	100	<u>CANOLA, LINOLA AND LINSEED</u> DO NOT apply after the 8 leaf stage of the crop. DO NOT apply after the commencement of stem elongation. This means that application must not occur after the 8 leaf stage, or if stem elongation commences before the 8 leaf stage, application must not occur after stem elongation has commenced. DO NOT apply more than 1 application of herbicide containing haloxyfop per crop. DO NOT apply after grazing. See GENERAL INSTRUCTIONS, Spraying oils/wetters section.
	Early Tillering	100	100	
Barley grass, Brome grass, Paradoxa grass, Volunteer cereals	2 to 4 leaf	50	75	<u>FIELD PEAS AND CANOLA:</u> The only oil recommended for use with EuroChem FEATHERTOP 520 is Uptake* Spraying Oil. EuroChem FEATHERTOP 520 + Lontrel*750 SC + Uptake* Spraying Oil are compatible and selective to canola. This tank-mix is also compatible with atrazine and selective to triazine tolerant canola.
	Early Tillering	75	100	
Wild oats WA, SA, Vic, Tas, Southern and Central NSW	2 to 4 leaf	37.5	50	<u>LUPINS AND FIELD PEAS:</u> Mixtures with Brodal* or simazine may cause crop yellowing and separate applications are recommended.
	Early Tillering	50	75	
Wild oats Northern NSW & Qld	2 to 4 leaf	50	75	<u>CHICKPEAS, FABA BEANS. LENTILS AND VETCH, LINOLA. LINSEED:</u> Broadleaf herbicides should not be added to EuroChem FEATHERTOP 520. Apply EuroChem FEATHERTOP 520 and broadleaf herbicides at least a week apart.
	Early Tillering	75	100	
				<u>LUCERNE. CLOVER OR MEDIC PASTURES:</u> If grazed or cut for hay immediately prior to treatment delay application until all grasses have fully expanded leaves. Use 75 mL + spraying oil or 100 mL + wetter/ha. (See GENERAL INSTRUCTIONS, Spraying Oils/wetters section). If silver grass (<i>Vulpia</i> spp.) is present in pasture, simazine should be tank mixed with the higher rate of EuroChem FEATHERTOP 520 plus a non-ionic wetter.

Table 1b. Winter crop growth stage application windows.

Crop	Crop growth Stage
Lucerne, Medic and Clover pastures or seed crops	Apply from 2 nd trifoliate leaf onwards. For <i>Erodium</i> spp. spraying, apply from cotyledon crop stage onwards.
Canola, Linola and Linseed	Apply from 2 leaf to 8 leaf stage of crop growth. DO NOT apply after the commencement of stem elongation. This means that application must not occur after the 8 leaf stage, or if stem elongation commences before the 8 leaf stage, application must not occur after stem elongation has commenced.
Chickpeas, Faba beans, Field peas, Lentils, Lupins, Vetch	Apply from 2 nd leaf, 2 nd node or 2 nd branch to prior to flowering.

Table 2a. Lucerne, Medic and Clover seed crops and pastures.

See table 1b for crop stages.

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE(mL/ha) with Uptake Spraying Oil	CRITICAL COMMENTS
Prairie grass (<i>Bromus catharticus</i>)	Up to early tillering	100	<u>See GENERAL INSTRUCTIONS, Spraying oils/wetters section.</u> ③Use lower rate when growing conditions and crop or pasture competition are good and when weed populations are below 100 plants/m². Use the higher rate when weed populations are above 100 plants/m² or when crop or pasture competition is poor. <u>NOTE:</u> Storksbill may not be controlled if simazine or Broadstrike* are tank-mixed with EuroChem FEATHERTOP 520. <u>LUCERNE, CLOVER OR MEDIC PASTURES:</u> If grazed or cut for hay immediately prior to treatment delay application until all grasses have fully expanded leaves. Use 75 mL + spraying oil or 100 mL + wetter/ha. (See GENERAL INSTRUCTIONS, Spraying Oils/wetters section). If silver grass (<i>Vulpia</i> spp.) is present in pasture, simazine should be tank mixed with the higher rate of EuroChem FEATHERTOP 520 plus a non-ionic wetter.
Musky or ferny leaf Storksbill (<i>Erodium moschatum</i>) Common Crowsfoot or Common Storksbill (<i>Erodium cicutarium</i>).	Up to 6 leaf or 5 cm diameter	50 - 75③	
Long or shiny leaf storksbill (<i>E. botrys</i>).	Up to 8 leaf or 5 cm diameter	75 - 100	

Table 2b. Lucerne, Medic and Clover seed crops only - not to be used for stockfeed.
See table 1b for crop stages.

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE(mL/ha) with Uptake Spraying Oil	CRITICAL COMMENTS
Couch grass (suppression), Rhodes grass (control)	Tillering seedlings	150 + 150④	④For best suppression of couch or control of Rhodes grass, make 2 applications of EuroChem FEATHERTOP 520 2 - 4 weeks apart. Time second application to coincide with tillering stage of weeds and just after irrigation or significant rain. Only treat actively growing weeds which are not moisture stressed. Use these rates for control of couch and Rhodes grass
Couch grass (control) Rhodes grass (control)	Established stands	400 - 800	

Table 3a. Summer crops - Cotton, Cowpea, Lucerne, Mung bean, Navy beans, Peanuts, Soybeans, Sunflowers.

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE(mL/ha) with Uptake Spraying Oil	CRITICAL COMMENTS
Australian millet	2 leaf to tillering up to 15 cm	150	See GENERAL INSTRUCTIONS, Spraying oils/wetters section.
Barnyard grass	2 to 5 leaf	100	
	Tillering up to 15 cm	150	<u>NAVY BEANS. PEANUTS. SOYBEANS:</u> For broadleaf weed control, EuroChem FEATHERTOP 520 at 150 mL/ha plus wetter may be tank mixed with Blazer* (except on navy beans) or Basagran*. Tank mixtures may cause transient leaf spotting on the crop but do not normally affect yield. DO NOT tank mix broadleaf herbicides with EuroChem FEATHERTOP 520 if grasses have begun tillering or if the grasses are under moisture stress. DO NOT add Uptake Spraying Oil when mixing with Blazer* or Basagran*. DO NOT use Blazer or Basagran* tank-mixes on Cowpeas.
Crowsfoot grass, Green panic, Johnson grass (rhizome)	2 leaf to tillering up to 15 cm	150	
Johnson grass (seedling), Liverseed grass (seedling), Mossman river grass	2 to 5 leaf	100	
	Tillering and up to 15cm	150	
Summer grass	2 leaf to tillering up to 15 cm	150	
Volunteer cereals	2 to 4 leaf	100	
	Tillering up to 15 cm	150	

Table 3b. Summer crop growth stage application windows.

Crop	Crop Growth Stage
Lucerne	Apply from 2 nd trifoliate leaf onwards.
Cowpea, Mung beans, Navy Beans, Soybeans	Apply from 2 nd leaf to flowering.
Peanuts	Apply from 2 nd leaf to pegging.
Cotton	Apply from 2 nd leaf to before the onset of flowering.
Sunflowers	Apply from 2 nd leaf to head initiation.

Table 4. Annual and Perennial grasses and *Erodium* spp. in Orchard, Vine and Plantation crops, forestry and pyrethrum.

CROPS	CROP GROWTH STAGE	WEEDS CONTROLLED	WEED GROWTH STAGE	RATE (mL/ha) with Uptake Spraying Oil	CRITICAL COMMENTS
Orchard, vine and plantation Crops including: Apples, Avocado, Banana, Blueberry, Citrus, Custard apple, Feijoa, Grapevines, Guava, Kiwifruit, Litchi (Lychee), Longan, Mango, Nashi, Nut trees, Passionfruit, Paw paw, Pear, Persimmon, Pineapple, Rambutan, Stone fruit.	All growth stages	<u>Perennial grasses:</u> Couch, Rhodes grass, Slender rats tail grass	Established stands	400 - 800	See GENERAL INSTRUCTIONS, Spraying oils/wetters section. Spray should be directed to the base of the tree or vine avoiding contact with fruit and foliage. Spot spray: Use 25 mL to 50 mL/100 L of water. Use higher rate on late tillering mature grasses. Annual Grasses: Where treated in association with perennial grasses, these annual grasses will be controlled. Forestry: For annual grasses apply lowest rate to newly emerged grasses, increasing the rate as they develop. Pyrethrum Tasmania only: For <i>Erodium</i> spp. apply 75 - 100 mL/ha if the main weed is <i>E. botrys</i>. Use 50 - 75 mL/ha if either <i>E. cicutarium</i> or <i>E. moschatum</i> are the main weeds.
		Buffel grass, Green panic, Johnson grass, Kikuyu, <i>Paspalum</i> spp, <i>Setaria</i> spp	Vegetative to early tillering	200	
			Late tillering	400	
		<u>Annual grasses:</u> Annual ryegrass, Barley grass, Barnyard grass, Brome grass, Crowsfoot grass, Lesser canary grass, Liverseed grass, Mossman river grass, Paradoxa grass, Summer grass, Volunteer cereals, Wild oats.	2 leaf to tillering	200	
Forestry. <i>Pinus radiata</i> <i>Eucalyptus</i> spp.					
Forestry: <i>Pinus pineaster</i>		Annual grasses as above	Vegetative to tillering	125 - 250	
Pyrethrum		Barley grass, Brome grass, Rope twitch, Barnyard grass, <i>Erodium</i> spp., Volunteer cereals.	Vegetative to tillering	100 - 250	

Table 5. EuroChem FEATHERTOP 520 and Select® Herbicide tank-mixes — Canola, Chickpeas, Faba beans Field peas, Lupins, Lentils.

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE (mL/ha)		CRITICAL COMMENTS
		EuroChem FEATHERTOP 520	Select Herbicide	
FOP/DIM susceptible Annual ryegrass + Volunteer barley, Volunteer wheat, Brome grass, Wild oats, Barley grass, Phalaris.	2 to 4 leaf	25	150	See GENERAL INSTRUCTIONS, Spraying oils/wetters section. Use Uptake* Spraying Oil at 500 mL/100 L or Hasten® at 1 L/100 L. Apply at the same crop growth stages as those in Table 1b Winter Crops. Lentils: Apply up to 7 node-early branching crop growth stage only. Lupins: Not for Qld.
	Early Tillering	38	150	
FOP resistant Annual ryegrass + Volunteer barley, Volunteer wheat, Brome grass, Wild oats, Barley grass, Phalaris.	2 to 4 leaf	25	200	
	Early Tillering	38	250	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

* = Registered Trademark

GENERAL INSTRUCTIONS

Mixing

- Add water to the spray tank to 10 cm above the level of agitation and ensure the agitation device is working vigorously. (There must be a minimum of 100 L of water in the tank before any pesticide is added.)
- If tank mixing, firstly, add any soluble liquid formulations (e.g. LONTREL* Herbicide) and allow agitation for approximately one minute.
- Then add EuroChem FEATHERTOP 520 at the point where agitation is strongest. (Do not add EuroChem FEATHERTOP 520 through a strainer or sieve). Allow further agitation for one minute.
- Half fill the spray tank.
- If using wettable powder or water dispersible granules, or other emulsifiable concentration formulations (e.g. LORSBAN* 750WG or Omethoate) these should be added after the EuroChem FEATHERTOP 520 to the half full spray tank ensuring vigorous agitation.
- Finally add Uptake* Spraying Oil or approved alternate spraying oil/wetter. (See *section on spraying oils/wetters*) and continue filling the tank to the required volume maintaining agitation at all times.
- Only mix sufficient solution for immediate use. EuroChem FEATHERTOP 520 and any other tank mixes should be applied immediately for best results.

Spraying Oils/wetters

Spraying Oils: It is essential to add an adjuvant to EuroChem FEATHERTOP 520. Best results will be achieved with Uptake* Spraying Oil at 0.5 L/100 L of spray solution. Alternatively, other oils plus a non-ionic wetter may also be used. When other crop spraying oils are used, mix at 1.0 L/100 L and add a *non-ionic wetter (surfactant)* at 200 mL/100 L of spray solution. Use of oil is not always recommended. See Critical Comments for specific situation recommendations.

Non-ionic Wetters: When Uptake or other oils are not used, a 100% concentrate non-ionic wetting agent such as BS-1000* at 200 mL/100 L must be used along with the higher rate of EuroChem FEATHERTOP 520 as specified in the Directions for Use.

Where water volumes of less than 50 L/ha are used, DO NOT use less than 250 mL/ha of Uptake or 500 mL/ha for oils other than Uptake or less than 100 mL/ha of wetter.

CANOLA, LUCERNE, MEDIC AND CLOVER PASTURES AND SEED CROPS:

When tank mixing EuroChem FEATHERTOP 520 with Lontrel herbicides (canola only) or Broadstrike (lucerne, clover and medics), use Uptake Spraying Oil with the lower rates of EuroChem FEATHERTOP 520 or a wetting agent with the higher rates of EuroChem FEATHERTOP 520 unless otherwise specified. When mixing EuroChem FEATHERTOP 520 with other broadleaf herbicides on these crops, DO NOT use an oil, use a wetter instead.

FIELD PEAS AND CANOLA:

The oil recommended is Uptake Spraying Oil. Hasten is also recommended for use with tank-mixtures of EuroChem FEATHERTOP 520 and Select Herbicide.

For canola, EuroChem FEATHERTOP 520 + Lontrel 750SG + Uptake Spraying Oil are compatible and selective to canola. This tank-mixture is also compatible with atrazine or simazine and selective to triazine tolerant canola.

NAVY BEANS, PEANUTS, SOYBEANS:

When mixing with Blazer or Basagran DO NOT add spraying oil to these mixtures. DO NOT use these tank-mixes on cowpea.

COMPATIBILITY

Ground use only: EuroChem FEATHERTOP 520 Herbicide can be tank mixed with:

Insecticides:	dimethoate Lorsban* 500 EC Insecticide Lorsban* 750WG Insecticide omethoate
Herbicides:	atrazine Basagran* Blazer* Broadstrike Herbicide Lontrel* Herbicide Lontrel* 750SG MCPA ester (LVE) - <i>DO NOT exceed 700 mL/ha of MCPA LVE</i> Oryzalin Select* Herbicide simazine Starane 200 Herbicide
Fungicides:	Dithane DF* Dithane Rainshield
Trace elements:	magnesium sulphate zinc sulphate

EuroChem FEATHERTOP 520 Herbicide is NOT COMPATIBLE with 2,4-D or MCPA as sodium or amine salts.

Aerial use: No product other than a recommended crop oil or wetter should be mixed with EuroChem FEATHERTOP 520 Herbicide when applied by air except for addition of Lontrel Forestry Herbicide for use in forestry and Lontrel 750SG for use in canola only.

APPLICATION

Apply EuroChem FEATHERTOP 520 Herbicide in sufficient water to obtain good coverage. It should be applied by an accurately calibrated ground rig or aircraft delivering droplets with a VMD of 200 - 300 microns.

The following spray volumes are recommended.

Ground application 50 - 150 L/ha
Aerial application 30 L/ha minimum

Use higher water volumes in orchards and in dense crops where the weeds may be shielded by the crop canopy.

CLEANING SPRAY EQUIPMENT

If broadleaf herbicides, particularly sulfonyleureas have been used in the spray equipment at any

time prior to EuroChem FEATHERTOP 520, particular care should be taken to follow the directions on the relevant broadleaf herbicide label for equipment cleaning, or damage to susceptible crops may occur.

After using EuroChem FEATHERTOP 520, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose, drain the tank and clean any filters in the tank, pump, line and nozzles.

To rinse. After cleaning the tank as above, quarter fill the tank with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

To decontaminate. Before spraying cereals, maize, sorghum or other sensitive crops, wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (e.g. SURF*, Cold Water SURF Concentrate*, DynamoMatic Concentrate*, OMO* or DRIVE*) at 500 mL/100 L of water or the powder equivalent at 500 g/100 L of water, and circulate throughout the system for at least fifteen minutes. Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain. Chlorine based cleaners are not recommended.

Rinse water should be discharged onto a designated disposal area, or if this is unavailable, onto unused land away from desirable plants and water sources.

* Registered trademarks